

# Palm kernel meal and cassava peel meal in growing pigs

Palm kernel meal and cassava peel meal in growing pigs

Growth performance and nutrient digestibility of growing pigs fed a mixture of palm kernel meal and cassava peel meal

## Abstract

In a 42 days feeding trials, twenty growing crossbred pigs with mean initial body weights of  $16 \pm 0.8$ kg were assigned randomly to 5 dietary treatment groups of four pigs each. A basal diet was formulated. The basal diet was based on cassava flour, maize offals, soyabean meal and brewer's dried grain (BDG). A mixture of cassava peel meal and (CPM) and palm kernel meal (PKM) replaced the BDG fraction in the basal diet in the ratio 2:0, 0:2, 1:1 and 1.75:1.75 in diets II, III, IV and V respectively. Each diet was offered on

*ad libitum*

basis. Pigs were individually weighed on weekly basis until the end of the experiment. Daily voluntary feed intake was monitored.

Growing pigs responded non-significantly ( $p > 0.05$ ) to the mixture of different fibre sources in body weight gain, feed conversion ratio and protein efficiency ratio except for feed intake, which varied significantly ( $p < 0.05$ ). Data on final body weight was described using the linear regression method. The slope of regression of final body weight (y) (kg/pig/week) depending on weeks (kg/week) was highest on diet V ( $3.525 \pm 0.1842$ ) and lowest on diet II ( $2.074 \pm 0.1742$ ). Digestibility values also varied significantly ( $p < 0.05$ ) with ether extract having the average digestibility value (%) of 90.32 and ash the lowest value of 53.60.

### Key words:

dietary fibre, additivity, efficiency of utilisation, weight gain



Department of Animal Science, University of Ibadan, Ibadan.

\*Department of Animal Science, Obafemi Awolowo University, Ile-Ife.

Yes