

Taenia solium control in Tanzania

Taenia solium control in Tanzania

Impact of farmer education on their knowledge, attitudes, and practices in southern Tanzania: a case for

Taenia solium

 control

Abstract

A comparative study was conducted with pre- and post-intervention assessments on the same subjects to evaluate a health education intervention administered to smallholder pig farmers in Iringa Rural (n = 750) and Chunya (n = 700) districts, southern Tanzania. A total of 366 respondents (249 in Iringa Rural, 117 in Chunya) undertook a self assessment questionnaire regarding knowledge on

Taenia solium

, its impact and mode of transmission, and attitudes towards consumption of infected pork.

McNemar and Bhapkar Chi-square P values were computed in SPSS and MH to assess significant differences.

The study revealed that health education intervention significantly improved the knowledge and attitudes towards

Taenia solium

 control (P < 0.001) and thus would reduce the consumption of infected pork. It

recommends that health education, particularly to women be integrated as an essential component of prevention and control programmes for

Taenia solium

 infections. The Government of Tanzania should revise its pork inspection regulations to include practical ways of controlling porcine cysticercosis.

Key words:

Health education effect, paired analysis, porcine cysticercosis control



H A Ngowi, E M Mkupasi, F P Lekule*, A L Willingham III and S M Thamsborg****

Department of Veterinary Medicine and Public Health, Sokoine University of Agriculture, P. O. Box 3021, Morogoro, Tanzania

** Department of Animal Science and Production, Sokoine University of Agriculture, P. O. Box 3004, Morogoro, Tanzania*

*** WHO/FAO Collaborating Center for Research and Training on Neglected and other Parasitic Zoonoses, Department of Veterinary Disease Biology, The Faculty of Life Sciences, University of Copenhagen, Dyrlægevej 100, 1870 Frederiksberg C, Denmark*

In partnership with [LRDD journal](#)

Yes