

Sugar cane molasses and cassava meal as alternative energy sources

Sugar cane molasses and cassava meal as alternative energy sources

Use of different protein levels in diets of sugar cane molasses type B or cassava meal and NUPROVIM based on soybean meal for growing-fattening pigs

Summary

Fifty six castrated male pigs, YL (Camborough) cross, of approximately 25 Kg live weight and 96 days average age were used. The pigs were allotted into a factorial arrangement (2x2) according to a random blocks design. The considered effects were energy source (sugar cane molasses type B or cassava meal) and the protein level in the diets (medium or low) given through the soybean meal.

For the energy source effect the dry matter intake was higher ($p < 0.001$) in pigs fed with the diet based on sugar cane molasses type B regarding to the diet based on cassava meal. To the level protein effect, there was increment in the dry matter intake ($p < 0.05$) and the protein conversion was significantly inferior ($p < 0.001$) when the protein low level was used; the daily gain was superior ($p < 0.001$) using protein medium level and also the alimentary conversion improved for this level.

It is considered the sugar cane molasses type B and cassava meal constitute alternative energy sources with the possibility to substitute totally to the cereals in the diets for growing-fattening

Keywords:

Cassava meal, growing-fattening pigs, performance, protein level, soybean meal, sugar cane molasses type B



C M Mederos*

S Rodríguez**

N Mazas**

R E Almaguel*

*Instituto de Investigaciones Porcinas, Gaveta Postal No. 1, Punta Brava, La Habana, Cuba

**Instituto Nacional de Investigaciones en Viandas Tropicales

with the partnership of

[Livestock Research and rural Development](#)

Oui