

# Parvovirus and PCV2 in crossbred Indian pigs

Parvovirus and PCV2 in crossbred Indian pigs

Porcine parvovirus- and porcine circovirus 2-associated reproductive failure and neonatal mortality in crossbred Indian pigs

*Rinku Sharma and G Saikumar*

*Tropical Animal Health and Production. Volume 42, Number 3 / mars 2010*

## Abstract

The objective of this study was to detect the presence of porcine parvovirus (PPV) and porcine circovirus 2 (PCV2) in a farm showing reproductive failure and increased mortality in neonatal piglets by histopathological examination, polymerase chain reaction, and demonstration of viral antigen and nucleic acid. Out of 594 piglets farrowed by 70 first-parity gilts, nine (1.51%) mummified fetuses, 13 (2.19%) stillborn, and 572 (96.3%) live-born piglets were recorded. The average litter size at birth was 8.48 piglets per litter. One hundred ninety-four (33.91%) piglets died within 7 days of age. PPV was detected in five litters (7.14%) and two of them revealed coinfection with PCV2. The pathological lesions in the coinfecting litters were more severe, indicating a synergistic action between the two viruses. Results of this study suggest for the first time occurrence of PPV and coinfection with PCV2 in crossbred Indian pigs affected with reproductive problem and neonatal mortality.

### **Keywords**

Pigs - Porcine parvovirus - Porcine circovirus 2 - Coinfection - India

Oui