

Palm kernel meal and cassava peel meal in growing pigs

Palm kernel meal and cassava peel meal in growing pigs

Growth performance and nutrient digestibility of growing pigs fed a mixture of palm kernel meal and cassava peel meal

Abstract

In a 42 days feeding trials, twenty growing crossbred pigs with mean initial body weights of 16 ± 0.8 kg were assigned randomly to 5 dietary treatment groups of four pigs each. A basal diet was formulated. The basal diet was based on cassava flour, maize offals, soyabean meal and brewer's dried grain (BDG). A mixture of cassava peel meal and (CPM) and palm kernel meal (PKM) replaced the BDG fraction in the basal diet in the ratio 2:0, 0:2, 1:1 and 1.75:1.75 in diets II, III, IV and V respectively. Each diet was offered on

ad libitum

basis. Pigs were individually weighed on weekly basis until the end of the experiment. Daily voluntary feed intake was monitored.

Growing pigs responded non-significantly ($p > 0.05$) to the mixture of different fibre sources in body weight gain, feed conversion ratio and protein efficiency ratio except for feed intake, which varied significantly ($p < 0.05$). Data on final body weight was described using the linear regression method. The slope of regression of final body weight (y) (kg/pig/week) depending on weeks (kg/week) was highest on diet V (3.525 ± 0.1842) and lowest on diet II (2.074 ± 0.1742). Digestibility values also varied significantly ($p < 0.05$) with ether extract having the average digestibility value (%) of 90.32 and ash the lowest value of 53.60.

Key words:

dietary fibre, additivity, efficiency of utilisation, weight gain



Department of Animal Science, University of Ibadan, Ibadan.

*Department of Animal Science, Obafemi Awolowo University, Ile-Ife.

Si